arXiv:gr-qc/9312028v1 17 Dec 1993

Distributional Energy-Momentum Tensor of the Kerr-Newman Space-Time Family

Herbert BALASIN [**] [*]
Institut für Theoretische Physik, Technische Universität Wien
Wiedner Hauptstraße 8–10, A - 1040 Wien, AUSTRIA

and

Herbert NACHBAGAUER [††]
Laboratoire de Physique Théorique ENSLAPP [†]
Chemin de Bellevue, BP 110, F - 74941 Annecy-le-Vieux Cedex, France

Abstract

Using the Kerr-Schild decomposition of the metric tensor that employs the algebraically special nature of the Kerr-Newman space-time family, we calculate the energy-momentum tensor. The latter turns out to be a well-defined tensor-distribution with disk-like support.

PACS numbers: 9760L, 0250

ENSLAPP-A-453/93
TUW 93 – 28
gr-qc/9312028

[**] e-mail: hbalasin @ email.tuwien.ac.at

[*] partially supported by the "Hochschuljubiläumsstiftung d. Stadt Wien" project H-00013

[††] e-mail: herby @ lapphp1.in2p3.fr

[†] URA 14-36 du CNRS, associée à l'E.N.S. de Lyon, et au L.A.P.P. (IN2P3-CNRS) d'Annecy-le-Vieux

Introduction

Singular space-times present one of the major challenges in general relativity. Originally it was believed that their singular nature is due to the high degree of symmetry of the well-known examples ranging from the Schwarzschild geometry to the Friedmann-Robertson-Walker cosmological models. However, Penrose and Hawking [1] have shown in their celebrated singularity theorems that singularities are a phenomenon which is inherent to general relativity. Since the standard approach allows only for smooth space-time metrics, one has to exclude the so-called singular regions from the space-time manifold.

In a recent work [2] the authors advocated the use of such distributional techniques to calculate the energy-momentum tensor of the Schwarzschild geometry. It turns out that it is possible to include the singular region (i.e. the space-like line $r = 0$ with respect to Schwarzschild coordinates) in the space-time which now no longer is a vacuum geometry, and to identify it with the support of the energy-momentum tensor. The latter becomes a tensor-distribution [3, 4] with delta-like shape.

This reasoning puts the Schwarzschild geometry on the same footing with its ultrarelativistic limit the Aichelburg-Sexl shock-wave geometry [5], where the energy-momentum tensor has a delta-like support on a null line and is interpreted as being generated by a particle moving with velocity of light. Adopting this line of arguments one might consider the Schwarzschild geometry as being generated by a tachyon which hides behind the event horizon thus providing a new interpretation of cosmic censorship.

The aim of the present work is to extend our calculation to the general axisymmetric, stationary space-time family discovered by Kerr and Newman [6, 7]. This family also contains the Schwarzschild geometry and its charged extension the Reissner-Nordstrøm solution as special cases of spherical symmetry. We will show that the distributional techniques developed in [2] apply to this family too thereby allowing to calculate its energy-momentum tensor.

One of the main features of the Kerr-geometry (and its charged version) which eventually led to its discovery is the existence of a geodetic, null vector-field k^a which gives rise to a corresponding congruence of geodesics. Taking advantage of this fact it is possible to decompose the metric into the so-called Kerr-Schild form [8], $g_{ab} = \eta_{ab} + f k_a k_b$, where η_{ab} denotes a flat (background) metric and f a scalar field. This geometrical decomposition greatly facilitates the calculation not only from a technical point of view but also from a conceptual one since it identifies k^a as being an integral part of the geometry which will be kept fixed during an eventual regularisation of g_{ab}.

Our work is organised in the following way: The first chapter is devoted to the calculation of the Ricci-tensor and the curvature scalar of an arbitrary metric of Kerr-Schild form. In the second chapter we will rederive our previous results concerning the Schwarzschild geometry and extend them to the Reissner-Nordstrøm case. Finally, in the third section we calculate the distributional energy-momentum tensor of the Kerr- and Kerr-Newman geometries.

1) Algebraically special geometries and Kerr-Schild structure

All geometries we are going to consider in this work are algebraically special geometries, which allow a Kerr-Schild decomposition of the metric

$$g_{ab} = \eta_{ab} + f k_a k_b, \tag{1}$$

where $k^a = \eta^{ab} k_b$ denotes a null vector (field) with respect to the metric η_{ab}, which in turn implies its nullity with respect to g_{ab}. The above decomposition provides two metrical structures associated with a given manifold $\mathcal{M}$, g_{ab} and η_{ab} respectively. In what follows ∇_a denotes the derivative operator associated with g_{ab} and ∂_a its commuting counterpart with respect to η_{ab}. An important consequence of the nullity of k^a and the decomposition (1) is

$$k^a \nabla_a k^b = k^a \partial_a k^b, \tag{2}$$

which can be derived from the explicit form of the difference operation ${C^a}_{bc}$ of the derivative operator

$$\begin{aligned}
&\nabla_a v^b = \partial_a v^b + {C^b}_{ad} v^d, \qquad v^a \in \Gamma(T\mathcal{M}), \qquad (3)\\
&{C^a}_{bc} = \frac{1}{2} g^{ad} \left(\partial_b g_{dc} + \partial_c g_{db} - \partial_d g_{bc}\right)\\
&\quad = \frac{1}{2} \left(\partial_b (f k^a k_c) + \partial_c (f k^a k_b) - \partial^a (f k_b k_c) + f k^a (k\partial)(f k_b k_c)\right),
\end{aligned}$$

where all index-raising and lowering is done with respect to η_{ab}. Equation (2) tells us that if k^a is geodetic with respect to g_{ab} the same is true with respect to η_{ab} and vice-versa. Kerr and Schild [8] have shown that the vacuum Einstein-(Maxwell)-field equations require k^a to be geodetic, and we can assume k^a to be affinely parametrised, i.e. $(k\partial)k^b = 0$. This condition is an essential property of the geometry and thus we will strictly maintain it, even in the course of a regularisation procedure.

The calculation of the Ricci-tensor using the conventions [9]

$$\begin{aligned}
&{R_{abd}}^c = \partial_b {C^c}_{ad} - \partial_a {C^c}_{bd} + {C^c}_{fb} {C^f}_{ad} - {C^c}_{fa} {C^f}_{bd},\\
&R_{ab} = {R_{acb}}^c = \partial_c {C^c}_{ab} - \partial_a {C^c}_{cb} + {C^c}_{cf} {C^f}_{ab} - {C^c}_{af} {C^f}_{cb},
\end{aligned}$$

together with the form (3) of ${C^a}_{bc}$, which implies ${C^a}_{ab} = 0$, yields

$$\begin{aligned}
&{R^a}_b = \frac{1}{2} \Big(\partial^a \partial_c (f k^c k_b) + \partial_b \partial^c (f k_c k^a) - \partial^2 (f k^a k_b)\Big), \qquad (4)\\
&R = \partial_a \partial_b (f k^a k^b).
\end{aligned}$$

Let us again remind the reader that from now on all indices are raised and lowered with the help of η_{ab}. (4) has the remarkable property of being a sum of second derivatives linear with respect to f which is neither the case for the upper nor the lower index parts. This property will allow a distributional evaluation of (4) whenever $f k^a k^b$ is itself a well-defined distribution and therefore possesses a natural second derivation.

2) Schwarzschild geometry and Reissner Nordstrøm extension

A simple illustration of the above concepts is provided by the Schwarzschild geometry and its Reissner-Nordstrøm extension. The Kerr-Schild form of the Schwarzschild geometry, whose line element reads in Schwarzschild coordinates

$$ds^2 = -dt^2\left(1-\frac{2m}{r}\right) + dr^2\left(1-\frac{2m}{r}\right)^{-1} + r^2 d\Omega^2,$$

is most easily displayed using the coordinate transformation

$$\bar{t} = t - 2m\log(2m-r), \qquad r < 2m$$

which gives

$$ds^2 = -d\bar{t}^2 + dr^2 + r^2 d\Omega^2 + \frac{2m}{r}(d\bar{t} - dr)^2 \tag{5}$$

and allows the immediate identification

$$f = \frac{2m}{r}, \quad k^a = (1, e^i_r), \qquad i = 1,2,3$$

where e^i_r is the unit vector with respect to η_{ab} of the spherical-coordinate system. Due to the stationarity of the metric (5) the distributional evaluation of the Ricci-tensor and the curvature-scalar reduce to a 3-dimensional problem on the $\bar{t} = const$ surfaces. Let us evaluate the curvature scalar explicitly acting on an arbitrary test-function $\varphi \in\ C_0^\infty(I\!R^3)$,

$$\begin{aligned}(R,\varphi) &= \lim_{\epsilon\to 0} \int\limits_{R^3-B_\epsilon} d^3x\, f k^i k^j\, \partial_i\partial_j\varphi(x) = \qquad (6)\\ &= \lim_{\epsilon\to 0}\Bigg[-\int\limits_{S_\epsilon} dA\, f k^i k^j\, N_i\partial_j\varphi(\epsilon e^i_r) + \int\limits_{S_\epsilon} dA\, \partial_i(f k^i k^j) N_j \varphi(\epsilon e^i_r) + \\ &\qquad + \int\limits_{R^3-B_\epsilon} d^3x\, \varphi(x)\partial_i\partial_j(f k^i k^j)\Bigg]\end{aligned}$$

where B_ϵ is the ϵ-ball around the origin and S_ϵ its boundary. $N^i = r^2 \sin\theta \, e_r^i$ is the outward directed normal of S_ϵ and dA denotes $d\theta d\phi$. The last term in (6) is just the curvature scalar which vanishes in the region $I\!R^3 - B_\epsilon$, the first term is already of order ϵ. Thus only the second term contributes and we get in the limit $\epsilon \longrightarrow 0$

$$(R, \varphi) = \lim_{\epsilon \longrightarrow 0} \int\limits_{S_\epsilon} dA \frac{2m}{\epsilon^2} \, \epsilon^2 \sin\theta \, \varphi(\epsilon e_r^i) = 8\pi m \, \varphi(0)$$

which is precisely the result we obtained in [2] in Schwarzschild coordinates. With respect to Kerr-Schild coordinates the energy-momentum tensor becomes

$$8\pi \, T^a{}_b = R^a{}_b - \frac{1}{2}\delta^a{}_b R = -8\pi m \, \delta^{(3)}(x)(\partial_{\bar{t}})^a \, (d\bar{t})_b.$$

The extension of this result to the Reissner-Nordstrøm metric is easily achieved by observing that we only have to replace

$$f = \frac{2m}{r} \quad \longrightarrow \quad f + \tilde{f} = \frac{2m}{r} - \frac{e^2}{r^2}.$$

Since expression (4) is linear in f we necessarily obtain an additional contribution $\tilde{R}^a{}_b$ to the Ricci-tensor. Note that $\tilde{f}$ is still locally integrable, which is a necessary condition for the existence of the distribution R. Let us exemplify the calculation of $\tilde{R}^a{}_b$ by the evaluation of $\tilde{R}^0{}_0$.

$$(\tilde{R}^0{}_0, \varphi) = \lim_{\epsilon \longrightarrow 0} \frac{1}{2} \int\limits_{R^3 - B_\epsilon} d^3x \, \tilde{f} \partial^2 \varphi =$$

$$\frac{1}{2} \lim_{\epsilon \longrightarrow 0} \left[- \int\limits_{S_\epsilon} dA \, \tilde{f} (N\partial) \varphi + \int\limits_{S_\epsilon} dA \, (N\partial) \tilde{f} \, \varphi + \int\limits_{R^3 - B_\epsilon} d^3x \, \partial^2 \tilde{f} \, \varphi \right] =$$

$$\frac{1}{2} \left[\int\limits_{S_\epsilon} dA \left(\frac{e^2}{\epsilon^2} \right) \epsilon^2 \sin\theta (e_r \partial) \varphi(0) + \int\limits_{S_\epsilon} dA \, \epsilon^2 \sin\theta \left(\frac{2e^2}{\epsilon^3} \right) (\varphi(0) + \epsilon (e_r \partial) \varphi(0)) \right.$$

$$\left. - \int\limits_{R^3 - B_\epsilon} d^3x \left(\frac{2e^2}{r^4} \right) (\varphi(x) - \varphi(0)) - \varphi(0) \int\limits_{R^3 - B_\epsilon} d^3x \, \left(\frac{2e^2}{r^4} \right) \right] =$$

$$- e^2 \int d^3x \, \frac{1}{r^4} (\varphi(x) - \varphi(0)) =: -(\left[\frac{e^2}{r^4} \right], \varphi),$$

where beginning with the third line of the calculation the limit $\epsilon \to 0$ is considered implicitly thereby dropping all terms of order ϵ. The calculation of the remaining components proceeds along the same lines so that we get in the end

$$\tilde{R}^0{}_0 = -\left[\frac{e^2}{r^4}\right], \quad \tilde{R}^0{}_i = 0, \quad \tilde{R}^i{}_j = \left[\frac{e^2}{r^4}(\delta^{ij} - 2e^i_r e^j_r)\right], \quad \tilde{R} = 0.$$

The electromagnetic part of the energy-momentum tensor is still traceless which tells us that the regularisation procedure did not destroy the conformal invariance of the theory.

With respect to Kerr-Schild coordinates the total energy-momentum tensor of the Reissner-Nordstrøm geometry is given by

$$T^0{}_0 = -m\delta^{(3)}(x) - \frac{e^2}{8\pi}\left[\frac{1}{r^4}\right], \quad T^0{}_i = 0, \quad T^i{}_j = \frac{e^2}{8\pi}\left[\frac{1}{r^4}(\delta^{ij} - 2e^i_r e^j_r)\right].$$

This result emphasizes the fact that the electromagnetic part of $T^a{}_b$ is a well-defined tensor-distribution, which coincides with the classical result for $r \neq 0$, i.e.

$$(\left[\frac{e^2}{r^4}\right], \varphi) = (\frac{e^2}{r^4}, \varphi)$$

for all $\varphi \in C_0^\infty(\mathbb{R}^3)$ that vanish at $r = 0$.

3) Kerr and Kerr-Newman geometries

Passing from Schwarzschild to Kerr generalises the null vector-field k^a and the function f to

$$\begin{aligned} k^a &= (1, k^i) = (1, \frac{rx + ay}{r^2 + a^2}, \frac{ry - ax}{r^2 + a^2}, \frac{z}{r}) \\ f &= \frac{2mr}{\Sigma}, \qquad \Sigma = \frac{r^4 + a^2 z^2}{r^2}, \end{aligned}$$

where r is implicitly given by

$$r^4 - r^2(x^2 + y^2 + z^2 - a^2) - a^2 z^2 = 0. \tag{7}$$

The cartesian coordinate system $\{x, y, z\}$ refers to the background metric η_{ab} of the Kerr-Schild decomposition. Taking again advantage of the stationarity of the metric reduces the distributional evaluation to a 3-dimensional problem. The constraint (7) may be solved by a change of coordinates from cartesian to spheroidal

$$\begin{aligned} x &= \sqrt{r^2 + a^2} \sin\theta \cos\phi \\ y &= \sqrt{r^2 + a^2} \sin\theta \sin\phi \\ z &= r \cos\theta. \end{aligned}$$

These coordinates represent a deformation of spherical coordinates expressed by the parameter a. The $r = const$ surfaces become confocal ellipsoids with focus on the ring $\rho^2 = x^2 + y^2 = a^2$, $z = 0$, whereas the $\theta = const$ surfaces are hyperboloids with an asymptotic cone of aperture θ. For $r = 0$ the ellipsoid degenerates into a double cover of the disk $\rho \leq a$, $z = 0$. In the latter region the coordinates essentially reduce to polar coordinates (ρ, ϕ) with radius $\rho = a \sin\theta$.

The calculation proceeds in a fairly straightforward fashion, using the general formulas of chapter one. This time, however, we have to exclude a disk-shaped region $r \leq \epsilon$ from the integrals and consider the limit $\epsilon \to 0$ afterwards in order to do partial integrations. To illustrate the procedure explicitly let us calculate the curvature-scalar given by (6) where N_i denotes the surface normal of the $r = const$ surface and dA the coordinate-surface area-element. Taking into account the identities

$$\begin{aligned} N^i &= r\sqrt{r^2 + a^2} \sin^2\theta e^i_\rho + (r^2 + a^2) \sin\theta \cos\theta e^i_z, \qquad (8) \\ k^i &= \frac{\sin\theta}{\sqrt{r^2 + a^2}} (r e^i_\rho - a e^i_\phi) + \cos\theta e^i_z, \quad (Nk) = \Sigma \sin\theta, \\ (N\partial)k^i &= \frac{a \sin^2\theta}{\sqrt{r^2 + a^2}} (a e^i_\rho + r e^i_\phi), \\ (k\partial)f + f(\partial k) &= \frac{2m}{\Sigma}, \end{aligned}$$

where e^i_ρ, e^i_ϕ and e^i_z denote the unit vectors of the cylindrical coordinates of $\mathbb{R}^3$, facilitates the evaluation of (6)

$$(R,\varphi) = \int\limits_{S_\epsilon} dA \frac{2m}{\Sigma}\, \Sigma \sin\theta\, \varphi(a\sin\theta\cos\phi, a\sin\theta\sin\phi, 0) =$$
$$\frac{4m}{a}\int\limits_0^{2\pi} d\phi \int\limits_0^{a} \frac{\rho d\rho}{\sqrt{a^2-\rho^2}} \varphi(\rho\cos\phi, \rho\sin\phi, 0)$$

which finally gives

$$R(x) = \frac{4m}{a}\frac{\vartheta(a-\rho)}{\sqrt{a^2-\rho^2}}\delta(z).$$

The calculation of the components of the Ricci-tensor proceeds along similar lines and gives

$$\begin{aligned}
{R^0}_0 &= 2m\left[\frac{a\vartheta(a-\rho)}{\sqrt{a^2-\rho^2}^3}\right]\delta(z) - \frac{2m}{a}\delta(\rho-a)\delta(z), \qquad (9)\\
{R^0}_i &= 2m\left[\frac{\rho\vartheta(a-\rho)}{\sqrt{a^2-\rho^2}^3}\right]\delta(z)e^i_\phi - \frac{m\pi}{a}\delta(\rho-a)\delta(z)e^i_\phi,\\
{R^i}_j &= -\frac{2m}{a}\left[\frac{\rho^2\vartheta(a-\rho)}{\sqrt{a^2-\rho^2}^3}\right]\delta(z)e^i_\phi e^j_\phi + \frac{2m}{a}\frac{\vartheta(a-\rho)}{\sqrt{a^2-\rho^2}}\delta(z)e^i_z e^j_z +\\
&\qquad \frac{4m}{a}\delta(\rho-a)\delta(z)e^i_\phi e^j_\phi
\end{aligned}$$

where

$$(\left[\frac{\vartheta(a-\rho)}{\sqrt{a^2-\rho^2}^3}\right]\delta(z),\varphi) :=$$
$$\int\limits_0^{a} d\rho \int\limits_0^{2\pi} d\phi \frac{\rho}{\sqrt{a^2-\rho^2}^3}\left(\varphi(\rho\cos\phi,\rho\sin\phi,0) - \varphi(a\cos\phi, a\sin\phi, 0)\right)$$

defines the distribution denoted by the square brackets. Our result shows that the energy distribution is concentrated on a disk with radius a in the $z=0$ plane. For all test functions that vanish on the circle $\rho = a$ our result coincides with that obtained in the classical work [10] on the source of the Kerr-geometry. Moreover, it can be shown that the limit $a \to 0$ exists and coincides with the result obtained in the Schwarzschild case. Let us demonstrate this fact for the curvature scalar

$$\lim_{a\to 0}(R_K,\varphi) = \lim_{a\to 0} 4m \int\limits_0^a d\rho \int\limits_0^{2\pi} d\phi \frac{\rho}{a\sqrt{a^2-\rho^2}} \varphi(\rho e^i_\rho, 0) =$$
$$8\pi m \int\limits_0^1 \frac{x dx}{\sqrt{1-x^2}}\, \varphi(0) = 8\pi m \varphi(0) = (R_S,\varphi),$$

where R_K and R_S denote the curvature scalars of the Kerr and the Schwarzschild geometry respectively.

Passing from Kerr to its electromagnetic extension the Kerr-Newman geometry merely changes the function

$$f = \frac{2mr}{\Sigma} \;\to\; f + \tilde{f} = \frac{2mr}{\Sigma} - \frac{e^2}{\Sigma}$$

in the Kerr-Schild decomposition (1), as it was the case with Schwarzschild and Reissner-Nordstrøm. The innocent looking additional contributions due to $\tilde{f}$ turn out to require considerable computational effort , although all integrals that are involved can be done analytically. In order to spare the reader the unwieldy formulas we will just give some useful identities needed for the calculation

$$\begin{aligned} &\partial_i k_j = \frac{r}{\Sigma} T_{ij} - \frac{a\cos\theta}{\Sigma}\epsilon_{ijk}k^k, \quad T_{ij} = \delta_{ij} - k_i k_j, \\ &(k\partial)\tilde{f} + (\partial k)\tilde{f} = 0, \quad (N\partial)\tilde{f} = \frac{2e^2(r^2+a^2)r\sin\theta}{\Sigma^2} \end{aligned} \tag{10}$$

where ϵ_{ijk} the standard ϵ-tensor of $\mathbb{R}^3$. Together with the identities (8), (10) leads to the following result for the $\tilde{f}$-contribution $\tilde{R}^a{}_b$ to the Ricci-tensor and to the curvature scalar

$$\tilde{R}^0{}_0 = -\frac{3\pi e^2}{4}\partial_i \left(\frac{\delta(\rho - a)}{a}\delta(z) e^i_\rho \right) - \left[\frac{e^2(r^2 + a^2 + a^2 \sin^2\theta)}{\Sigma^3} \right], \tag{11}$$

$$\begin{aligned}
\tilde{R}^0{}_i = &-\frac{e^2}{a}\partial_k \left(\frac{\vartheta(a-\rho)}{\sqrt{a^2-\rho^2}}\delta(z)(e^k_\rho e^i_\rho + e^k_\phi e^i_\phi) \right) - \frac{3\pi e^2}{4}\partial_k \left(\frac{\delta(\rho - a)}{a}\delta(z) e^k_\rho e^i_\phi \right) \\
&- \frac{\pi e^2}{2a}\left(\frac{\delta(\rho - a)}{a}\delta(z)(e^i_\rho - e^i_\phi) \right) + e^2 \left[\frac{\rho}{a}\frac{\vartheta(a-\rho)}{\sqrt{a^2-\rho^2}^3}\delta(z) e^i_\rho \right] \\
&- 2e^2 \left[\frac{a \sin\theta}{\Sigma^3}\sqrt{r^2 + a^2} e^i_\phi \right],
\end{aligned}$$

$$\begin{aligned}
\tilde{R}^i{}_j = &\frac{2e^2}{a}\partial_k \left(\frac{\rho}{a}\frac{\vartheta(a-\rho)}{\sqrt{a^2-\rho^2}}\delta(z)(e^k_\rho e^{(j}_\rho e^{i)}_\phi + e^k_\phi e^{(j}_\phi e^{i)}_\phi \right) \\
&+ \frac{3\pi e^2}{4}\partial_k \left(\frac{\delta(\rho - a)}{a}\delta(z) e^i_\phi e^j_\phi e^k_\rho \right) - \frac{2e^2}{a^2}\left[\frac{\rho^2 \vartheta(a-\rho)}{\sqrt{a^2-\rho^2}^3}\delta(z) \right] e^{(i}_\rho e^{j)}_\phi \\
&+ \frac{4e^2}{a}\frac{\delta(\rho - a)}{a}\delta(z) e^{(i}_\rho e^{j)}_\phi + \left[\frac{e^2}{\Sigma^3}((r^2 + a^2)\cos^2\theta - r^2 \sin^2\theta)(e^i_\rho e^j_\rho - e^i_z e^j_z) \right] \\
&+ \left[\frac{e^2}{\Sigma^3}(r^2 + a^2 + a^2 \sin^2\theta) e^i_\phi e^j_\phi \right] - \frac{4e^2}{\Sigma^3} r\sqrt{r^2 + a^2}\sin\theta\cos\theta e^{(i}_\rho e^{j)}_z,
\end{aligned}$$

$$\tilde{R} = \frac{2e^2}{a}\partial_k \left(\frac{\rho}{a}\frac{\vartheta(a-\rho)}{\sqrt{a^2-\rho^2}}\delta(z) e^k_\phi \right),$$

where the parenthesis around the indices of the base vectors denotes symmetrisation with unit weight. The non-vanishing trace of the Ricci-tensor is due to the length-scale a of the disk. However, conformal invariance is restored in the limit $a \to 0$, thus reproducing the result of the spherical symmetric case.

Let us finally comment on the issue of regularisation. Our calculation started from the Kerr-Schild decomposition of the metric. In the derivation of the Ricci-tensor and the curvature-scalar we implicitly assumed the validity of classical differential calculus and the existence of a smooth function f. This might be interpreted as regularising the intermediate quantities like the difference tensor of the covariant derivatives. Since the final result, the Ricci-tensor, turned out to be the second derivation of a distribution it was possible to directly evaluate it without referring to any regularisation procedure.

Conclusion

In the present work we have explicitly shown how to calculate the distributional energy-momentum tensor of the Kerr-Newman space-time family, thereby following closely the approach proposed in [2] which includes the singular regions of the geometry in the manifold. The Kerr-Schild-structure related to the algebraically special nature of this space-time family turned out to play a prominent role of the interpretation. Moreover, our results furnish a well-defined basis for the investigation of the so-called ultrarelativistic limit geometries, [5, 11] by boosting the energy-momentum tensor. Work in this direction is currently under progress.

Acknowledgement: The authors are greatly indebted to Prof. P. C. Aichelburg for many useful discussions.

References

[1] Hawking S W and Ellis G F R, *The large scale structure of space-time* Cambridge University Press, 1973.

[2] Balasin H and Nachbagauer H, *Class. Quantum Grav.* **10**, 2271 (1993).

[3] Lichnerowicz A, *Propagateurs, Commutateurs et Anticommutateurs en Relativité Générale*, IHES No. 10 (1961).

[4] Parker P E, Distributional Geometry, *J. Math. Phys.* **20** (1979) 1423.

[5] Aichelburg P and Sexl R, *J. Gen. Rel. Grav.* **2** (1971) 303.

[6] Kerr R P, *Phys. Rev. Lett.* **11** (1963) 237;
Newman E T and Janis A I, *J. Math. Phys.* **6** (1965) 915.

[7] Boyer R H and Lindquist R W, *J. Math. Phys.* **8** (1967) 265.

[8] Debney G C, Kerr R P and Schild A, *J. Math. Phys.* **10** (1969) 183.

[9] Wald R, *General Relativity*, University of Chicago Press, 1984.

[10] Israel W, *Phys. Rev.* **D 2** (1970) 641; Hamity V H, *Phys. Lett.* **A56** (1976) 77.

[11] Loustó C O and Sánchez N, *Nucl. Phys.* **B383** (1992) 377.